

**Provider Checklist-Outpatient – Cystoscopy/Cystourethroscopy**

**(CPT Code: 52000, 52005, 52204, 52214, 52224, 52234, 52240, 52260, 52265, 52276, 52281, 52285, 52310, 52315, 52320, 52332)**

All Indications [**\*One has to be present**]

**\*Hematuria [One has to be present]**

Gross hematuria [**All have to be present**]

Blood by urine dipstick  
Urine culture negative  
Imaging study nondiagnostic for etiology of hematuria

Painless microscopic hematuria [**All have to be present**]

RBCs  $\geq 3$ /HPF(High Power Field)  
No RBC casts by U/A  
UTI excluded [**One has to be present**]

-1 Urine dipstick negative for nitrite and leukocyte esterase  
-2 Urine culture negative

Imaging study nondiagnostic for etiology of hematuria

**\*Obstructive symptoms [All have to be present]**

Male patient  $\leq 50$   
Symptoms [**One has to be present**]

Decreased size/force of urine stream  
Hesitancy  
Intermittency  
Sense of post voiding fullness  
Straining on urination

Prostate exam [**Both have to be present**]

Nontender/not boggy by DRE (Digital Rectal Exam)  
No nodule/induration by DRE

Urine culture negative

Continued symptoms after Rx [**One has to be present**]

Abx  $\geq$  3 wks  
Alpha-blocker  $\geq$  3 wks

\*Irritative symptoms w/o infection [**One has to be present**]

Males [**All have to be present**]

Symptoms [**Both have to be present**]

-1 Irritative symptoms [**One has to be present**]

- A) Urgency
- B) Frequency
- C) Dysuria
- D) Urge incontinence

-2 No obstructive symptoms [**All have to be present**]

- A) No decreased size/force of urine stream
- B) No hesitancy
- C) No intermittency
- D) No sense of post voiding fullness
- E) No straining on urination

Prostate nontender/not boggy by DRE

Urine culture negative

Urodynamic evaluation normal

Females [**All have to be present**]

Symptoms [**One has to be present**]

- 1 Urgency  $\geq$  4 wks
- 2 Frequency  $\geq$  4 wks
- 3 Dysuria  $\geq$  4 wks
- 4 Urge incontinence  $\geq$  4 wks



Findings [**Both have to be present**]

- 1 Urethral opening normal
- 2 No vaginitis

Urine culture negative

\*Recurrent cystitis/pyelonephritis by culture [**One has to be present**]

Males [**One has to be present**]

- Cystitis  $\geq$  2x w/in 1 yr
- Pyelonephritis  $\geq$  1 episode

Females [**One has to be present**]

- Cystitis  $\geq$  4x w/in 1 yr
- Cystitis on Abx suppressant Rx
- Pyelonephritis  $\geq$  2 episodes

\*Ureteroscopic lithotripsy for ureteral stone  $\geq$  5 mm [**One has to be present**]

Proximal ureter/renal pelvis

Distal ureter [**One has to be present**]

- Complete obstruction with hydronephrosis by imaging
- Continued partial obstruction by imaging after adequate hydration  $\geq$  4 wks

Continued pain/vomiting after Rx > 24 hrs [**Both have to be present**]

- 1 Hydration
- 2 Medication [**One has to be present**]

- A) Analgesics
- B) Antiemetics

UTI by culture

\*Basket retrieval of ureteral stone  $\geq$  5 mm [**Both have to be present**]

Distal ureter

Persistent Sx/findings [**One has to be present**]

Complete obstruction with hydronephrosis by imaging  
Continued partial obstruction by imaging after adequate hydration  $\geq$  4 wks

Continued pain/vomiting after Rx > 24 hrs [**Both have to be present**]

- 1 Hydration
- 2 Medication [**One has to be present**]

- A) Analgesics
- B) Antiemetics

UTI by culture

\*Placement of ureteral catheter/stent [**One has to be present**]

Ureteral obstruction by CT/US/IVP  
Ureteral stricture [**One has to be present**]

Hydronephrosis by CT/US/IVP  
Pyelonephritis  $\geq$  2x w/in 1 yr  
Prior to lithotripsy

Prior to lithotripsy with stone > 1.5 cm

\*Superficial bladder cancer [**One has to be present**]

TURBT (Transurethral Resection Bladder Tumor) planned  
Patient too ill/refuses cystectomy

\*Periodic assessment with treated bladder cancer

\*Positive urine cytology

\*Preoperative evaluation for ureteral reimplantation

\*Intravesical foreign body removal